

**AQA, Edexcel, OCR**

**A Level**

# **A Level Physics**

## **Gravitational Fields 1**

Name:

**M M E**

Mathsmadeeasy.co.uk

Total Marks: /30

1.

Total for Question 1: 13

(a) Which two of the following statements are false? [2]

- i. Any object with mass will generate a gravitational field.
- ii. Gravitational fields require two masses.
- iii. A gravitational field is one of numerous fields that give rise to forces.
- iv. The strength of a planet's gravitational field is inversely proportional to the square of the distance from the surface.

(b) Describe what happens to the gravitational force between two objects A and B when:  
i. Their separation doubles. [1]

ii. The mass of A halves. [1]

iii. The mass of A doubles and that of B halves. [1]

iv. The mass of B triples and the separation halves. [2]

- (c) Sketch, for each of the following, the pattern of field lines. For the first three, by distributing your field lines accordingly, make the relative field strengths clear. [6]
- i. A sphere of mass  $m$ .
  - ii. A sphere of mass  $M$ , where  $m < M$ .
  - iii. A point source of mass  $M$ .
  - iv. A small section of a planet's surface.

2. Zog is the only planet in its solar system. It has a radius of 150 km and is perfectly spherical.

Total for Question 2: 13

- (a) By considering Newton's Second Law and his Law of Gravitation, derive an expression for the gravitational field strength,  $g$ , of an object in terms of its mass,  $m$ , the distance from its centre of mass,  $r$ , and the gravitational constant,  $G$ . [2]

- (b) Kyle measures a gravitational acceleration of  $0.5 \text{ ms}^{-2}$  when his spaceship is 1.0 km from Zog's surface. Calculate the average density of Zog. [4]

Though Zog is small, it has an even smaller moon, whose radius is 5 km. The separation of their centres of masses is 200 km. The resultant gravitational field is zero at a distance of 40 km from Zog's surface.

(c) By equating the gravitational field strengths, calculate the mass of the moon.


[4]

(d) Calculate the resultant field halfway between their surfaces, specifying in which direction the resultant force of attraction acts.

[3]

3. The graph below shows how the measured gravitational field strength ( $\text{ms}^{-2}$ ) varied with  $1/r^2$  ( $r$  in m) in an experiment carried out by Zoe

Total for Question 3: 4



(a) Use the graph to calculate the mass of the object used.

[4]