

OCR

A Level

A Level Physics

Oscillations

Name:


M M E

Mathsmadeeasy.co.uk

Total Marks: /30

1. The graph below shows the displacement of a spring, x , against the time since its release, t .

Total for Question 1: 6



- (a) On the plot above, sketch graphs of velocity versus time and acceleration versus time. Hence conclude whether or not the spring is in simple harmonic motion. Justify your conclusion. [3]

- (b) The period of a pendulum in simple harmonic motion is independent of its amplitude. How would you verify this statement experimentally? [3]

2. Geoff, a rock climber, is following his friend Marie up an overhanging cliff. Marie is attached to the rock and ensures that the rope between herself and Geoff is taut at all times. Unfortunately, Geoff falls off and subsequently swings. Unless otherwise stated, ignore air resistance.

Total for Question 2: 24

- (a) Geoff passes through the equilibrium position, for the second time, after 4.5 s. At this point, he is travelling at a speed of 5 ms^{-1} . Calculate the following:

i. The frequency of his oscillations.

[2]

ii. The amplitude of his oscillations.

[1]

iii. His velocity 2.25 s after he falls.

[3]

For a simple pendulum undergoing simple harmonic motion, $\omega^2 = \frac{g}{l}$, where l is the length of the pendulum and g is the acceleration due to gravity. For the small angles involved in simple harmonic motion, $\cos \theta \approx 1 - \frac{\theta^2}{2}$ and $\sin \theta \approx \theta$.

- (b) By expressing the angle θ in terms of Geoff's displacement, x , and the rope's length, l , derive an expression for $\cos \omega t$ in terms of l , A and θ . [2]

- (c) Using this result, show that his kinetic energy is given by $\frac{1}{2}m\omega^2(A^2 - l^2\theta^2)$. [3]

(d) By considering his vertical displacement, derive an expression for his potential energy in terms of l , g , θ and his mass, m . [3]

(e) Hence show that energy is conserved. [3]

(f) Sketch, on a single set of axes, the variation of E_{total} , E_p and E_k with θ for $-\theta_{max} \leq \theta \leq \theta_{max}$. [2]

- (g) Sketch, on a single set of axes, the displacements of a lightly damped and a very heavily damped simple pendulum as a function of time. In reality, after numerous oscillations, Geoff comes to rest. Is the damping in the system light, heavy or very heavy? [3]

- (h) In this case, the damping causes an exponential decrease in the amplitude of his displacements. Given that after 4 s the amplitude has reduced to 4.00 m, calculate the amplitude when $t = 8$ s. Use your answer to part a,ii as an initial amplitude. [2]