

**AQA**

**A Level**

# **A Level Physics**

**ELECTRICAL CIRCUITS: Electrical  
Quantities (Answers)**

Name:

**M M E**

**Mathsmadeeasy.co.uk**

**Total Marks: /30**

1. This question is about the variation of quantities such as current, voltage and resistance in simple electrical circuits containing a variety of standard components.

Total for Question 1: 11




Figure 1: I-V characteristics for three different circuit components.

- (a) State Ohm's Law. [1]

**Solution:**  $I \propto V$

- (b) Assign one of the following components to each of the characteristic graphs in Figure 1: filament lamp, semiconductor diode, resistor. [3]

**Solution:**  
1: resistor  
2: filament lamp  
3: diode

- (c) Why have these been plotted on graphs of current against potential difference rather than current against electromotive force? [1]

**Solution:** EMF is concerned with energy being put into the circuit; PD is concerned with energy used by components.

- (d) For the diode, state the value of the resistance when a backward bias is applied. [1]

**Solution:** Zero

(e) Sketch the following graphs:

[2]

- i. Resistance against temperature for an ntc thermistor.
- ii. Current against voltage for an ntc thermistor.

**Solution:**

- (i) Non-linear decrease of resistance as temperature increases. Concave up.
- (ii) inverse of the filament lamp i.e. concave up in the positive quadrant and concave down in the negative quadrant.

(f) The current in a filament is 8 A. In the time during which Patrick is using the lamp,  $8 \times 10^{22}$  electrons pass through a given point in the circuit. For how long has he been using the lamp?

[3]

**Solution:** 1600 s

2. James unexpectedly finds an electrical circuit in his physics classroom. Immediately he starts recording the current. He notes that it decreases linearly from 10 A to zero over a time period of 30 s.

Total for Question 2: 5

- (a) Plot a graph of current against time.

[2]

**Solution:** As described.

- (b) Calculate the charge that is transferred in this time.

[2]

**Solution:** 150 C

- (c) If James had also been able to record a graph of charge (vertical axis) against time (horizontal axis), which of the following accurately describes what he would have seen?

[1]

- i. Linear increase.
- ii. Non-linear increase.
- iii. Linear decrease.

**Solution:** Option 2.

3. Frances is exploring the electrical properties of a piece of wire. She observes that:
- (a) for a given current, doubling the length,  $L$ , of the wire doubles the potential difference (P.D.) and the resistance,  $R$ .
  - (b) for a given P.D., doubling the wire's diameter,  $d$ , causes  $R$  to decrease by a factor of 4.

Total for Question 3: 11

- (a) On the basis of Frances' observations, which of these relationships is true: [3]
- i.  $R \propto A$  and  $R \propto L$
  - ii.  $R \propto 1/A$  and  $R \propto 1/L$
  - iii.  $R \propto 1/A$  and  $R \propto L$
  - iv.  $R \propto d^2$  and  $R \propto L$

**Solution:** Option 3.

- (b) Use this to define resistivity,  $\rho$ , in terms of  $d$ ,  $R$  and  $L$ . [2]

**Solution:**  $\frac{\pi d^2 R}{4L}$

- (c) Figure 2 is a characteristic graph for a circuit component. Calculate the resistivity at the point for which the curves tangent has been drawn given that the component is cylindrical, has a length of 8 cm and has a radius of  $1.5 \times 10^{-5}$  m. [3]

**Solution:**  $1.8 \times 10^{-8}$




Figure 2: Characteristic graph for a particular circuit component.

- (d) Explain how, using the characteristic, it is possible to deduce that, for this component, resistivity increases with temperature. [3]

**Solution:** As temperature increases, so too does current. But, on the characteristic, as current increases the gradient decreases i.e.  $R$  increases. Since  $\rho = \frac{AR}{L}$ , if  $R$  increases but the dimensions remain unchanged,  $\rho$  will also increase.

4. This question is about superconductors.

Total for Question 4: 3

(a) A superconductor is a material whose resistance...

[1]

- i. ... increases to  $\infty$  below a specific critical temperature.
- ii. ... decreases to zero above a specific critical temperature.
- iii. ... decreases to zero below a specific critical temperature.

**Solution:** Option 3.

(b) At present the highest known critical temperature is approximately  $-130^{\circ}\text{C}$ . Give two examples that illustrate why a superconductor with a room temperature critical temperature would be particularly useful.

[2]

**Solution:** Any valid examples e.g. long-lasting batteries, heat-free laptops.