GCE Examinations

Advanced Subsidiary / Advanced Level

Further Pure Mathematics
Module FP2

Paper G

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.

Written by Rosemary Smith & Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

FP2 Paper G – Marking Guide

1. *(a)* $y = e^{\arctan x} \;\therefore\; \dfrac{dy}{dx} = \dfrac{1}{1+x^2} \times e^{\arctan x} = \dfrac{e^{\arctan x}}{1+x^2}$ M1 A1

$\dfrac{d^2y}{dx^2} = \dfrac{1}{1+x^2} \times \dfrac{e^{\arctan x}}{1+x^2} - (1+x^2)^{-2} \times 2x \times e^{\arctan x} = \dfrac{e^{\arctan x}(1-2x)}{(1+x^2)^2}$ M1 A1

(b) pt. of inflexion $\therefore \dfrac{d^2y}{dx^2} = 0$ M1

$\therefore e^{\arctan x} = 0$ (no solns) or $x = \frac{1}{2}$ A1

$\therefore$ coordinates (0.5, 1.59) [y-coord 3sf] A1 **(7)**

---

2. *(a)* let $y = \operatorname{arcosh} x \;\therefore \cosh y = x$

$\therefore \sinh y \dfrac{dy}{dx} = 1$ M1

$\dfrac{dy}{dx} = \dfrac{1}{\sqrt{\cosh^2 y - 1}} = \dfrac{1}{\sqrt{x^2-1}}$ M1 A1

(b) $u = \operatorname{arcosh} x,\ u' = \dfrac{1}{\sqrt{x^2-1}};\ v' = 1,\ v = x$ M1

$\displaystyle\int \operatorname{arcosh} x \; dx = x \operatorname{arcosh} x - \int \frac{x}{\sqrt{x^2-1}} \; dx$ A1

$= x \operatorname{arcosh} x - \sqrt{x^2-1} + c$ M1 A1 **(7)**

---

3. $t = \tan x \;\therefore\; \dfrac{dt}{dx} = \sec^2 x = (1 + t^2)$ M1 A1

$\displaystyle\int_0^{\frac{\pi}{4}} \frac{1}{1+\sin 2x} \; dx = \int_0^1 \frac{1}{1+\frac{2t}{1+t^2}} \times \frac{1}{1+t^2} \; dt$ M1 A1

$\displaystyle = \int_0^1 \frac{1}{1+t^2+2t} \; dt$ M1

$\displaystyle = \int_0^1 \frac{1}{(t+1)^2} \; dt$ A1

$= [-(t+1)^{-1}]_0^1$ A1

$= -\frac{1}{2} - (-1) = \frac{1}{2}$ A1 **(8)**

© *Solomon Press*

4. *(a)* $4x^2 - 4x + 10 \equiv (2x-1)^2 - 1 + 10 \equiv (2x-1)^2 + 9$ M1 A1

$$\int \frac{1}{\sqrt{4x^2-4x+10}}\,\mathrm{d}x = \int \frac{1}{\sqrt{(2x-1)^2+9}}\,\mathrm{d}x$$

$u = 2x - 1,\ \dfrac{\mathrm{d}u}{\mathrm{d}x} = 2$ M1

$= \frac{1}{2}\displaystyle\int \frac{1}{\sqrt{u^2+9}}\,\mathrm{d}u$ A1

$= \frac{1}{2}\operatorname{arsinh}\left(\frac{u}{3}\right) + c = \frac{1}{2}\operatorname{arsinh}\left(\frac{2x-1}{3}\right) + c$ M1 A1

(b) $\displaystyle\int_{\frac{1}{2}}^{2} \frac{1}{\sqrt{4x^2-4x+10}}\,\mathrm{d}x = \left[\frac{1}{2}\operatorname{arsinh}\left(\frac{2x-1}{3}\right)\right]_{\frac{1}{2}}^{2}$

$= \frac{1}{2}\operatorname{arsinh} 1 - \frac{1}{2}\operatorname{arsinh} 0$ M1

$= \frac{1}{2}\ln(1+\sqrt{2})$ M1 A1 **(9)**

---

5. *(a)*



B3

B2

(b) $2 - \tanh x = 3\ \operatorname{sech} x$

$2 - \dfrac{e^x - e^{-x}}{e^x + e^{-x}} = \dfrac{6}{e^x + e^{-x}}$ M1

$2e^x + 2e^{-x} - e^x + e^{-x} = 6$

$e^x + 3e^{-x} - 6 = 0$ A1

$e^{2x} - 6e^x + 3 = 0$ M1

$e^x = \dfrac{6 \pm \sqrt{24}}{2} = 3 \pm \sqrt{6}$ M1 A1

$\therefore x = \ln(3 \pm \sqrt{6}) = -0.60$ or 1.70 (2dp) M1 A1 **(12)**

© *Solomon Press*

6. *(a)* $u = \sin^{n-1}x,\ u' = (n-1)\sin^{n-2}x\cos x;\ \ v' = \sin x,\ v = -\cos x$ M1 A1

$I_n = [-\cos x \sin^{n-1}x]_0^{\frac{\pi}{2}} + \int_0^{\frac{\pi}{2}} (n-1)\sin^{n-2}x\cos^2 x\ \mathrm{d}x$ A1

$I_n = 0 - 0 + (n-1)\int_0^{\frac{\pi}{2}} \sin^{n-2}x(1-\sin^2 x)\ \mathrm{d}x$ M1 A1

$I_n = (n-1)\int_0^{\frac{\pi}{2}} \sin^{n-2}x\ \mathrm{d}x - (n-1)\int_0^{\frac{\pi}{2}} \sin^n x\ \mathrm{d}x$

$I_n = (n-1)I_{n-2} - (n-1)I_n$ M1

$nI_n = (n-1)I_{n-2}$

$I_n = \frac{n-1}{n} I_{n-2}$ A1

(b) curve sym. about $x = \frac{\pi}{2}\ \ \therefore V = \int_0^{\pi} \pi y^2\ \mathrm{d}x = 2\pi\int_0^{\frac{\pi}{2}} y^2\ \mathrm{d}x = 2\pi I_4$ M1 A1

$I_0 = \int_0^{\frac{\pi}{2}} \mathrm{d}x = [x]_0^{\frac{\pi}{2}} = \frac{\pi}{2}$ M1 A1

$I_2 = \frac{1}{2}I_0 = \frac{\pi}{4}$ M1

$I_4 = \frac{3}{4}I_2 = \frac{3\pi}{16}$ A1

$\therefore V = 4\pi \times \frac{3\pi}{16} = \frac{3}{4}\pi^2$ A1 **(14)**

---

7. *(a)* $x = a\cosh t,\ \frac{\mathrm{d}x}{\mathrm{d}t} = a\sinh t,\ \frac{\mathrm{d}^2x}{\mathrm{d}t^2} = a\cosh t$ M1 A1

$y = 2a\sinh t,\ \frac{\mathrm{d}y}{\mathrm{d}t} = 2a\cosh t,\ \frac{\mathrm{d}^2y}{\mathrm{d}t^2} = 2a\sinh t$ A1

at vertex $y = 0\ \therefore t = 0\ \ \therefore \frac{\mathrm{d}x}{\mathrm{d}t} = 0,\ \frac{\mathrm{d}^2x}{\mathrm{d}t^2} = a,\ \frac{\mathrm{d}y}{\mathrm{d}t} = 2a,\ \frac{\mathrm{d}^2y}{\mathrm{d}t^2} = 0$ A1

$\rho = \left|\frac{(0+(2a)^2)^{\frac{3}{2}}}{0\times 0 - 2a\times a}\right| = \left|\frac{8a^3}{-2a^2}\right| = 4a$ M1 A1

(b) $\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}t} \div \frac{\mathrm{d}x}{\mathrm{d}t} = \frac{2a\cosh t}{a\sinh t} = \frac{2\cosh t}{\sinh t}$ M1 A1

eqn. is $y - 2a\sinh p = \frac{2\cosh p}{\sinh p}(x - a\cosh p)$ M1

$y\sinh p - 2a\sinh^2 p = 2x\cosh p - 2a\cosh^2 p$

giving $2x\cosh p - y\sinh p = 2a$ A1

(c) asymptotes are "$y = \pm\frac{b}{a}x$" $\ \therefore\ y = \pm\frac{2a}{a}x = \pm 2x$ M1

$\therefore l_1$ is $y = 2x$; tangent at P is $2x\cosh p - y\sinh p = 2a$

$\therefore$ at $Q\ \ 2x\cosh p - 2x\sinh p = 2a$ M1

giving $x = \frac{a}{\cosh p - \sinh p}$ A1

"$b^2 = a^2(e^2-1)$" $\ \therefore 4a^2 = a^2(e^2-1),\ e > 0\ \ \therefore e = \sqrt{5}\ \ \therefore S$ is $(a\sqrt{5}, 0)$ M1 A1

$\therefore\ a\sqrt{5} = \frac{a}{\cosh p - \sinh p}\ \ \therefore \cosh p - \sinh p = \frac{1}{\sqrt{5}}$ M1

$\frac{1}{2}(\mathrm{e}^p + \mathrm{e}^{-p}) - \frac{1}{2}(\mathrm{e}^p - \mathrm{e}^{-p}) = \frac{1}{\sqrt{5}}$ M1

$\therefore\ \mathrm{e}^{-p} = \frac{1}{\sqrt{5}},\ -p = \ln 5^{-\frac{1}{2}},\ p = \frac{1}{2}\ln 5$ A1 **(18)**

Total **(75)**

© Solomon Press

Performance Record – FP2 Paper G

Question no.	**1**	**2**	**3**	**4**	**5**	**6**	**7**	Total
Topic(s)	diff. inv. trig.	diff., integr., inv. hyp. fns.	integr. $t = \tan x$	integr. std. forms	eqn. in hyp. fns.	reduction formula	hyperbola, rad. of curv., tangent	
Marks	7	7	8	9	12	14	18	75
Student								

© *Solomon Press*