



**General Certificate of Education (A-level)  
June 2012**

**Mathematics**

**MS03**

**(Specification 6360)**

**Statistics 3**

***Mark Scheme***

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: [aqa.org.uk](http://aqa.org.uk)

Copyright © 2012 AQA and its licensors. All rights reserved.

**Copyright**

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

## Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

## No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

## MS03

Q	Solution	Marks	Total	Comments
1(a)	$r = 0.665$ to $0.666$ $r = 0.6$ to $0.7$  <b>Alternative</b> Use of a correct formula  $r = 0.665$ to $0.666$	B2 (B1)	2	AWFW (0.66577) AWFW
(b)	$H_0: \rho = 0$ $H_1: \rho > 0$  SL $\alpha = 0.01$ (1%) CV $r = (+)0.658$  Calculated $r = 0.666 >$ Tabulated $r$  <b>Evidence</b> , at 1% level, of a <b>positive correlation</b> between the neck length and the tail length of mature male giraffes	B1          A1F	4	Both; do not accept in terms of $r$ but accept in words  AWRT (0.6581) $H_1: \rho \neq 0 \Rightarrow CV = (\pm)0.7079$  Comparison  F on $r$ and CV At 1% level, <b>reject hypothesis of no correlation</b>
<b>Total</b>			<b>6</b>	
2(a)	$H_0: \mu_F = \mu_M$ $H_1: \mu_F \neq \mu_M$  SL $\alpha = 0.01$ (1%) CV $z = 2.57$ to $2.58$  $z = \frac{\bar{f} - \bar{m}}{\sqrt{\frac{s_F^2}{n_F} + \frac{s_M^2}{n_M}}} = \frac{22.0 - 21.6}{\sqrt{\frac{1.31^2}{50} + \frac{0.702^2}{75}}}$  <b>1.95 to 2(.0)</b>  <b>No evidence</b> , at 1% level, to suggest that <b>there is a difference</b> between mean lengths	B1          A1F	6	Both; allow suffices of 1 & 2 or X & Y  AWFW (2.5758)  Numerator  Denominator; allow $\div 49$ and $\div 74$ but <b>not</b> pooling  AWFW (1.98)
(b)	Diameter, thickness, girth, width, weight, shape, colour, texture	B1	1	Accept any sensible alternative but <b>not</b> 'quality'
<b>Total</b>			<b>7</b>	

## MS03 (cont)

Q	Solution	Marks	Total	Comments
3(a)(i)	$P(S \cap U) = 0.15 \times 0.10 = \mathbf{0.015}$	B1	1	CAO
(ii)	$P(O \cap \geq 2) = (0.40 \times 0.50) + (0.45 \times 0.40)$ $+ (0.15 \times 0.70)$ $= 0.20 + 0.18 + 0.105 = \mathbf{0.485}$	M1 A1	2	$\geq 1$ term correct; may be implied CAO
(iii)	$P(U) = (0.40 \times 0.15) + (0.45 \times 0.05)$ $+ (0.15 \times 0.10)$ or (i) $= 0.06 + 0.0225 + 0.015 = \mathbf{0.097 \text{ to } 0.098}$	M1 A1	2	$\geq 2$ terms correct; may be implied AWFW (0.0975)
(iv)	$P(D   U) = \frac{P(D \cap U)}{P(U)} = \frac{0.40 \times 0.15}{\text{(iii)}}$ $= \frac{0.06}{0.0975} = \mathbf{0.612 \text{ to } 0.619}$	M1 A1	2	May be implied AWFW (0.61538)
(v)	$P(S   O) = \frac{0.15 \times (1 - 0.10)}{1 - \text{(iii)}} = \frac{0.135}{0.9025}$ $= \mathbf{0.149 \text{ to } 0.15}$	M1 M1 A1	3	Numerator Denominator AWFW (0.14958)
(b)	$P(D \cap T \cap S   O)$ $= \frac{0.40 \times 0.85}{1 - \text{(iii)}} \times \frac{0.45 \times 0.95}{1 - \text{(iii)}} \times \text{(v)} \times 3!$ $= \frac{0.34 \times 0.4275 \times 0.135 \times 6}{0.9025^3}$ <b>or</b> $= 0.3767 \times 0.4737 \times 0.1496 \times 6$ $= \mathbf{0.16}$	M1 M1 M1 A1	4	$\geq 2$ terms correct in numerator (1 - (iii)) in denominator 3! or 6 or 3 AWRT (0.16016)
<b>Total</b>			<b>14</b>	

## MS03 (cont)

Q	Solution	Marks	Total	Comments
4	$H_0: \lambda = 2.6$ (650) $H_1: \lambda > 2.6$ (650)	B1		Both; accept $\mu$ instead of $\lambda$
	SL $\alpha = 0.05$ (5%)			
	CV $z = 1.64$ to <b>1.65</b>	B1		AWFW (1.6449) Allow 1.96 iff $H_1: \lambda \neq 2.6$
	$\hat{\lambda} = \frac{688}{250} = 2.75(2)$	B1		AWRT (2.752) Can be implied by use of 688
	$z = \frac{2.752 - 2.6}{\sqrt{\frac{2.6}{250}}} = \frac{688 - 650}{\sqrt{650}} = 1.47$ to <b>1.49</b>	M1 A1		Allow use of 2.752 or 688 or 687.5 in denominator AWFW
	$p\text{-value} = 0.068$ to <b>0.071</b> > 0.05	(M1)		Use of $P(X \geq 688   \lambda = 650) = 0.072$ $\Rightarrow$ M0 A0 (M1) AF1
	<b>No evidence</b> , at 5% level, <b>to support</b> manager's suspicion	A1F	6	F on CV and $z$ -value
	<b>Total</b>		<b>6</b>	
5(a)	$\hat{p} = \frac{68}{125} = 0.544$	B1		CAO; or equivalent
	98% $\Rightarrow z = 2.32$ to <b>2.33</b>	B1		AWFW (2.3263)
	Approximate CI for $p$ is	M1		
	$0.544 \pm 2.3263 \times \sqrt{\frac{0.544 \times 0.456}{125}}$	A1F		F on $\hat{p}$ and $z$
	<b><math>0.544 \pm (0.103</math> to <b>0.104)</b></b>	A1		CAO/AWFW May be implied by <b>correct</b> answer AWFW
or				
<b><math>(0.44</math> to <b>0.441, 0.647</b> to <b>0.65)</b></b>	A1			
<b><math>(44\%, 65\%)</math></b>	A1	6	AWRT	
(b)	Require $2 \times 2.3263 \times \sqrt{\frac{p(1-p)}{n}} \leq 0.1$ (10%)	M1 A1		Allow 'no 2' and FT on CI from (a) Allow $p = 0.44$ to $0.65$
	Thus $\sqrt{n} = 22.1$ to $23.3$	M1		Attempted solution for $\sqrt{n}$ or $n$
	$n = 490, 495, \dots, 545$	A1	4	Must be to 'nearest 5'
	<b>Total</b>		<b>10</b>	

## MS03 (cont)

Q	Solution	Marks	Total	Comments
6(a)(i)	$M = U + V$ $E(M) = 13 + 15 = \mathbf{28}$	B1		CAO
	$V(M) = 3^2 + 6^2$ $+ 2 \times 3 \times 6 \times (-0.6)$ $= 9 + 36 - 21.6 = \mathbf{23.4}$	M1 M1 A1	4	Allow 'no 2' CAO
	(ii) $D = W - 2U$ $E(D) = 24 - 2 \times 13 = \mathbf{-2}$	B1		CAO
	$V(D) = 4^2 + (2^2 \times 3^2)$ $= 16 + 36 = \mathbf{52}$	M1 A1	3	CAO
(iii)	$T = M + W + X$ $E(T) = 28 + 24 + 9 = \mathbf{61}$	B1F		F on $E(M)$ from (a)(i)
	$V(T) = 23.4 + 4^2 + 2^2 = \mathbf{43.4}$	B1F	2	F on $V(M)$ from (a)(i)
(b)(i)	$P(M = 30) = \mathbf{0}$	B1	1	CAO
(ii)	$P(W > 2U) = P(D > 0) = P\left(Z > \frac{0 - (-2)}{\sqrt{52}}\right)$	M1		Standardising 0 using c's $E(D)$ & c's $V(D)$ from (a)(ii)
	$= P(Z > 0.28) = 1 - 0.61026$	m1		Area change
	$= \mathbf{0.389 \text{ to } 0.394}$	A1	3	AWFW (0.39076)
(iii)	$P(50 < T < 70) =$ $P\left(Z < \frac{70 - 61}{\sqrt{43.4}}\right) - P\left(Z < \frac{50 - 61}{\sqrt{43.4}}\right)$	M1		Standardising 70 & 50 using c's $E(T)$ & c's $V(T)$ from (a)(iii)
	$= P(Z < \mathbf{1.37}) - P(Z < \mathbf{-1.67})$	A1		AWRT either
	$= 0.91466 - (1 - 0.95254)$	m1		Area change
	$= \mathbf{0.865 \text{ to } 0.87}$	A1	4	AWFW (0.86657)
<b>Total</b>			<b>17</b>	

## MS03 (cont)

Q	Solution	Marks	Total	Comments
7(a)(i)	$X \sim B(n, p)$			
	$E(X) = \sum_{x=0}^n x \times \binom{n}{x} p^x (1-p)^{n-x}$	M1		Used; ignore limits until A1
	$= np \times \sum_{x=1}^{n-1} \frac{(n-1)!}{(x-1)!(n-x)!} p^{x-1} (1-p)^{n-x}$	M1		Factor of $np$ plus $p^x$ to $p^{x-1}$ , $n!$ to $(n-1)!$ and $x!$ to $(x-1)!$ Ignore limits until A1
	$= np \times \sum_{x=1}^{n-1} B(n-1, p) = np$	A1	3	Fully correct and complete derivation AG
(ii)	$\text{Var}(X) = E(X^2) - [E(X)]^2$	M1		Used
	$= E[X(X-1)] + E(X) - [E(X)]^2$ $= n(n-1)p^2 + np - n^2p^2 = np(1-p)$	A1	2	(Other derivations are possible) Fully correct and complete derivation
(b)(i)	$\binom{n}{m} p^m (1-p)^{n-m} \geq \binom{n}{m-1} p^{m-1} (1-p)^{n-m+1}$	M1		Stated or used
	$\frac{n!}{m!(n-m)!} p^{n-m} \geq \frac{n!}{(m-1)!(n-m+1)!} (1-p)^{n-m+1}$	m1		Expanding $\binom{n}{m}$ and $\binom{n}{m-1}$
	$(n-m+1)p \geq m(1-p)$ $np - mp + p \geq m - mp$	A1		Simplification
	$m \leq (n+1)p$	A1	4	Fully correct and complete derivation AG
(ii)	$B(10, 0.65)$			
	$(n+1)p = 7.15$ and $(n+1)p - 1 = 6.15$	M1		Can be implied by '7'
	$m = 7$	A1		CAO
	$B(35, 0.5)$			
	$(n+1)p = 18$ and $(n+1)p - 1 = 17$	(M1)		Can be implied by '17 or 18'
	$m = 17$ and $18$	A1	3	CAO; accept 'or'
(c)	$Y \sim B(4000, 0.00095) \approx \text{Po}(3.8)$	B1		May be implied
	$k = 3$	B1		CAO; may be implied
	$P(Y \leq 3) = 0.473$ to $0.474$	B1	3	AWFW (0.4735)
	SC: $k = 4 \Rightarrow P(Y \leq 4) = 0.667$ to $0.668$ Award B1 B0 B1			AWFW (0.6678)
	<b>Total</b>		<b>15</b>	
	<b>TOTAL</b>		<b>75</b>	