

Write your name here

Surname

Other names

**Pearson Edexcel**  
**Level 3 GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

**Mathematics**  
**Advanced Subsidiary**  
**Paper 1: Pure Mathematics**

Wednesday 16 May 2018 – Morning  
**Time: 2 hours**

Paper Reference

**8MA0/01**

**You must have:**

Mathematical Formulae and Statistical Tables, calculator

Total Marks

--

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

**Instructions**

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

**Information**

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 15 questions in this question paper. The total mark for this paper is 100.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

**Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P58346A

©2018 Pearson Education Ltd.

1/1/1/1/1/



  
**Pearson**

Answer ALL questions. Write your answers in the spaces provided.

1. Find

$$\int \left( \frac{2}{3}x^3 - 6\sqrt{x} + 1 \right) dx$$

giving your answer in its simplest form.

$$\int \left( \frac{2}{3}x^3 - 6\sqrt{x} + 1 \right) dx$$

(4)

$$= \int \frac{2}{3}x^3 dx - 6 \int \sqrt{x} dx + \int 1 dx$$

$$= \frac{2}{3} \cdot \frac{1}{4} x^4 - 6 \cdot \frac{2}{3} \cdot x^{3/2} + x + c$$

$$= \frac{1}{6} x^4 - 4x^{3/2} + x + c$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





2. (i) Show that  $x^2 - 8x + 17 > 0$  for all real values of  $x$

(3)

(ii) "If I add 3 to a number and square the sum, the result is greater than the square of the original number."

State, giving a reason, if the above statement is always true, sometimes true or never true.

(2)

$$(i) \quad x^2 - 8x + 17 = (x-4)^2 - 16 + 17 \\ = (x-4)^2 + 1$$

$(x-4)^2 + 1 > 0$ , because ~~the~~ when you square a number the answer is always  $\geq 0$ , so  $(x-4)^2 + 1 > 0$  for all real values of  $x$ .

$$(ii) \quad (x+3)^2$$

The statement is only sometimes true.

e.g. let  $x = 5$ ,  $(x+3)^2 = 64$ ,  $x^2 = 25$  (statement true)

let  $x = -5$ ,  $(x+3)^2 = 4$ ,  $x^2 = 25$ . (statement not true).

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



3. Given that the point  $A$  has position vector  $4\mathbf{i} - 5\mathbf{j}$  and the point  $B$  has position vector  $-5\mathbf{i} - 2\mathbf{j}$ ,

(a) find the vector  $\vec{AB}$ ,

(2)

(b) find  $|\vec{AB}|$ .

Give your answer as a simplified surd.

(2)

$$\begin{aligned} \text{a) } \vec{AB} &= \cancel{(4\mathbf{i} - 5\mathbf{j})} - \cancel{(5\mathbf{i} - 2\mathbf{j})} \\ &= (5\mathbf{i} - 2\mathbf{j}) - (4\mathbf{i} - 5\mathbf{j}) \\ &= -9\mathbf{i} + 3\mathbf{j} \end{aligned}$$

$$\text{b) } |\vec{AB}| = \sqrt{(-9)^2 + 3^2} = \sqrt{90} = 3\sqrt{10}.$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





4. The line  $l_1$  has equation  $4y - 3x = 10$   $y = \frac{3}{4}x + \frac{5}{2}$

The line  $l_2$  passes through the points  $(5, -1)$  and  $(-1, 8)$ .

Determine, giving full reasons for your answer, whether lines  $l_1$  and  $l_2$  are parallel, perpendicular or neither.

$$\text{gradient of } l_2 = \frac{8+5}{-1+1} = \frac{-1-8}{5+1} = -\frac{3}{2} \quad (+)$$

$$\text{gradient of } l_1 = \frac{3}{4}$$

$$\frac{3}{4} \times -\frac{3}{2} = -\frac{9}{8} \neq -1 \text{ so not } \perp$$

$$\frac{3}{4} \neq -\frac{3}{2} \text{ so not parallel.}$$

Hence neither.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



5. A student's attempt to solve the equation  $2\log_2 x - \log_2 \sqrt{x} = 3$  is shown below.

$$2\log_2 x - \log_2 \sqrt{x} = 3$$

$$2\log_2 \left( \frac{x}{\sqrt{x}} \right) = 3$$

using the subtraction law for logs

$$2\log_2 (\sqrt{x}) = 3$$

simplifying

$$\log_2 x = 3$$

using the power law for logs

$$x = 3^2 = 9$$

using the definition of a log

- (a) Identify two errors made by this student, giving a brief explanation of each.

(2)

- (b) Write out the correct solution.

(3)

a) 1<sup>st</sup> error : when using the subtraction law  
you need to accommodate for the 2, in  $2\log_2 x$ .

2<sup>nd</sup> error : if  $\log_2 x = 3$ , then  $x = 2^3 = 8$ .

$$b) \quad 2\log_2 x - \log_2 \sqrt{x} = 3$$

$$\Rightarrow \log_2 x^2 - \log_2 \sqrt{x} = 3$$

$$\Rightarrow \log_2 \left( \frac{x^2}{\sqrt{x}} \right) = 3$$

$$\Rightarrow \log_2 (x^{3/2}) = 3$$

$$\Rightarrow \frac{3}{2} \log_2 (x) = 3$$

$$\Rightarrow \log_2 (x) = 3 \times \frac{2}{3} = 2$$

$$\Rightarrow x = 2^2 = 4.$$





6.




Figure 1

A company makes a particular type of children's toy.

The annual profit made by the company is modelled by the equation

$$P = 100 - 6.25(x - 9)^2$$

where  $P$  is the profit measured in thousands of pounds and  $x$  is the selling price of the toy in pounds.

A sketch of  $P$  against  $x$  is shown in Figure 1.

Using the model,

- (a) explain why £15 is not a sensible selling price for the toy. (2)

Given that the company made an annual profit of more than £80 000

- (b) find, according to the model, the least possible selling price for the toy. (3)

The company wishes to maximise its annual profit.

State, according to the model,

- (c) (i) the maximum possible annual profit,  
 (ii) the selling price of the toy that maximises the annual profit. (2)

$$a) \quad x = 15, \quad P = 100 - 6.25(15 - 9)^2 = -125$$

So it isn't suitable as the company would make a loss.



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 6 continued

$$b) \quad 100 - 6.25(x-9)^2 > 80 \text{ } \cancel{000}$$

$$\Rightarrow 100 - 80 \text{ } \cancel{000} > 6.25(x-9)^2$$

$$\Rightarrow 20 > 6.25(x-9)^2$$

$$\Rightarrow 3.2 > (x-9)^2$$

$$\Rightarrow \pm\sqrt{3.2} > x-9$$

$$\Rightarrow x < \pm\sqrt{3.2} + 9$$

$$\text{and } x > -\sqrt{3.2} + 9$$

So the minimum price is £7.22

$$c) \text{ i) Maximum profit} = \text{£}100\,000$$

$$\text{(ii) Selling price} = \text{£}9.$$



P 5 8 3 4 6 A 0 1 3 4 8



7. In a triangle  $ABC$ , side  $AB$  has length 10 cm, side  $AC$  has length 5 cm, and angle  $BAC = \theta$  where  $\theta$  is measured in degrees. The area of triangle  $ABC$  is  $15 \text{ cm}^2$


(a) Find the two possible values of  $\cos \theta$

(4)

Given that  $BC$  is the longest side of the triangle,

(b) find the exact length of  $BC$ .

(2)



$$\text{Area} = 15$$

$$\text{a) Area} = \frac{1}{2} ab \sin C$$

$$\Rightarrow \frac{1}{2} \times 10 \times 5 \times \sin \theta = 15$$

$$\Rightarrow \sin \theta = \frac{3}{5}$$

$$\Rightarrow \theta = \sin^{-1}\left(\frac{3}{5}\right) = 36.87$$

$$\cos^2 \theta = 1 - \sin^2 \theta$$

$$\cos^2 \theta = 1 - \frac{9}{25}$$

$$\cos \theta = \pm \frac{4}{5}$$

$$\text{b) } c^2 = a^2 + b^2 - 2ab \cos C$$

$$BC^2 = 10^2 + 5^2 - 2 \times 10 \times 5 \cos \left( \sin^{-1}\left(\frac{3}{5}\right) \right)$$

$$BC = \sqrt{205}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



8. A lorry is driven between London and Newcastle.

In a simple model, the cost of the journey £ $C$  when the lorry is driven at a steady speed of  $v$  kilometres per hour is

$$C = \frac{1500}{v} + \frac{2v}{11} + 60$$

(a) Find, according to this model,

(i) the value of  $v$  that minimises the cost of the journey,

(ii) the minimum cost of the journey.

(Solutions based entirely on graphical or numerical methods are not acceptable.)

(6)

(b) Prove by using  $\frac{d^2C}{dv^2}$  that the cost is minimised at the speed found in (a)(i).

(2)

(c) State one limitation of this model.

(1)

$$a) \quad i) \quad C = \frac{1500}{v} + \frac{2v}{11} + 60$$

$$\frac{dC}{dv} = -\frac{1500}{v^2} + \frac{2}{11} + 60$$

$$\text{let } \frac{dC}{dv} = 0 = -\frac{1500}{v^2} + \frac{2}{11}$$

$$\Rightarrow \frac{2}{11} = \frac{1500}{v^2}$$

$$\Rightarrow v^2 = 8250, \quad v = 90.8 \text{ km h}^{-1}$$

$$(ii) \quad \text{let } v = 90.8, \quad C = \frac{1500}{90.8} + \frac{2(90.8)}{11} + 60$$

$$\text{min cost} = \pounds 93.03$$





Question 8 continued

$$b) \frac{d^2C}{dv^2} = \frac{3000}{v^3}$$

$$v = 90.8, \quad \frac{d^2C}{dv^2} = 0.004 > 0 \text{ hence min cost.}$$

c) It would be impossible to drive at this speed for the entire journey

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 8 3 4 6 A 0 1 9 4 8

9.

$$g(x) = 4x^3 - 12x^2 - 15x + 50$$

(a) Use the factor theorem to show that  $(x + 2)$  is a factor of  $g(x)$ .

(2)

(b) Hence show that  $g(x)$  can be written in the form  $g(x) = (x + 2)(ax + b)^2$ , where  $a$  and  $b$  are integers to be found.

(4)




Figure 2

Figure 2 shows a sketch of part of the curve with equation  $y = g(x)$

(c) Use your answer to part (b), and the sketch, to deduce the values of  $x$  for which

(i)  $g(x) \leq 0$

(ii)  $g(2x) = 0$

(3)

$$\begin{aligned} \text{a) let } x = -2, \quad g(-2) &= 4(-2)^3 - 12(-2)^2 - 15(-2) + 50 \\ &= -32 - 48 + 30 + 50 \\ &= 0 \end{aligned}$$

Hence  $(x+2)$  is a factor.

$$\begin{aligned} \text{b) } 4x^3 - 12x^2 - 15x + 50 &= (x+2)(4x^2 - 20x + 25) \\ &= (x+2)(2x-5)^2 \end{aligned}$$

$$\text{c) i) } g(-2) = 0, \text{ so } x \leq -2.$$

$$\text{(ii) } g(2x) = 0, \quad x = -1, \quad x = 1.25$$





10. Prove, from first principles, that the derivative of  $x^3$  is  $3x^2$

(4)

$$\text{let } y = x^3$$

$$\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h}$$

$$= \lim_{h \rightarrow 0} \frac{x^3 + 3x^2h + 3xh^2 + h^3 - x^3}{h}$$

$$= \lim_{h \rightarrow 0} \frac{3x^2h + 3xh^2 + h^3}{h}$$

$$= \lim_{h \rightarrow 0} 3x^2 + 3xh + h^2$$

$$= 3x^2 \text{ as } h \rightarrow 0.$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



11. (a) Find the first 3 terms, in ascending powers of  $x$ , of the binomial expansion of

$$\left(2 - \frac{x}{16}\right)^9$$

giving each term in its simplest form.

(4)

$$f(x) = (a + bx)\left(2 - \frac{x}{16}\right)^9, \text{ where } a \text{ and } b \text{ are constants}$$

Given that the first two terms, in ascending powers of  $x$ , in the series expansion of  $f(x)$  are 128 and  $36x$ ,

(b) find the value of  $a$ ,

(2)

(c) find the value of  $b$ .

(2)

$$\begin{aligned} \text{a) } \left(2 - \frac{x}{16}\right)^9 &= 2^9 + \binom{9}{1} 2^8 \cdot \left(-\frac{x}{16}\right) + \binom{9}{2} 2^7 \left(-\frac{x}{16}\right)^2 + \dots \\ &= 512 + 9 \cdot 2^8 \cdot \left(-\frac{x}{16}\right) + 36 \cdot 2^7 \left(-\frac{x}{16}\right)^2 + \dots \\ &= 512 - 144x + 18x^2. \end{aligned}$$

$$\begin{aligned} \text{b) } 128 &= 2^9 \cdot a \\ a &= \frac{1}{4} \end{aligned}$$

$$\begin{aligned} \text{c) } 36x &= 512bx - 144xa \\ 72 &= 512b \\ b &= \frac{9}{64}. \end{aligned}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





12. (a) Show that the equation

$$4\cos\theta - 1 = 2\sin\theta\tan\theta$$

can be written in the form

$$6\cos^2\theta - \cos\theta - 2 = 0 \quad (4)$$

(b) Hence solve, for  $0 \leq x < 90^\circ$

$$4\cos 3x - 1 = 2\sin 3x \tan 3x$$

giving your answers, where appropriate, to one decimal place.

(Solutions based entirely on graphical or numerical methods are not acceptable.)

(4)

$$a) \quad 4\cos\theta - 1 = 2\sin\theta\tan\theta$$

$$4\cos\theta - 1 = \frac{2\sin\theta \sin\theta}{\cos\theta}$$

$$4\cos\theta - 1 = \frac{2\sin^2\theta}{\cos\theta}$$

$$4\cos^2\theta - \cos\theta = 2\sin^2\theta$$

$$4\cos^2\theta - \cos\theta = 2(1 - \cos^2\theta)$$

$$4\cos^2\theta - \cos\theta = 2 - 2\cos^2\theta$$

$$6\cos^2\theta - \cos\theta - 2 = 0$$

$$b) \quad 4\cos 3x - 1 = 2\sin 3x \tan 3x$$

$$\Rightarrow 6\cos^2 3x - \cos 3x - 2 = 0$$

$$\Rightarrow (3\cos^2 3x - 2)(2\cos 3x + 1)$$

$$\cos 3x = \frac{2}{3} \quad \text{or} \quad \cos 3x = -\frac{1}{2}$$

$$x = 16.06 \quad \text{or} \quad 40, 80$$



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

13.




Figure 3

The value of a rare painting, £ $V$ , is modelled by the equation  $V = pq^t$ , where  $p$  and  $q$  are constants and  $t$  is the number of years since the value of the painting was first recorded on 1st January 1980.

The line  $l$  shown in Figure 3 illustrates the linear relationship between  $t$  and  $\log_{10} V$  since 1st January 1980.

The equation of line  $l$  is  $\log_{10} V = 0.05t + 4.8$

- (a) Find, to 4 significant figures, the value of  $p$  and the value of  $q$ . (4)
- (b) With reference to the model interpret
- the value of the constant  $p$ ,
  - the value of the constant  $q$ . (2)
- (c) Find the value of the painting, as predicted by the model, on 1st January 2010, giving your answer to the nearest hundred thousand pounds. (2)

$$a) \quad V = pq^t, \quad \log_{10} V = 0.05t + 4.8$$

$$\Rightarrow \log_{10} pq^t = 0.05t + 4.8$$

$$\text{If we let } t=0, \quad \log_{10} p = 4.8$$

$$\Rightarrow 4.8^{10} 10^{4.8} = p = 63095 = 63100 \text{ to 4 s.f.}$$

$$\text{let } t=1, \quad \log_{10} 10^{4.8} \cdot q = 0.05t + 4.8$$

$$\Rightarrow \log_{10} 10^{4.8} + \log_{10} q = 4.85$$

$$\Rightarrow \log_{10} q = 0.05$$

$$\Rightarrow q = 10^{0.05} = 1.122$$





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 13 continued

b) i) The value of the initial value of the painting.

(ii) The proportional increase of the value of the painting.

$$c) V = 10^{4.8} \times (10^{0.05})^{30} = \pounds 2\,000\,000$$



14. The circle  $C$  has equation

$$x^2 + y^2 - 6x + 10y + 9 = 0$$

(a) Find

(i) the coordinates of the centre of  $C$

(ii) the radius of  $C$

(3)

The line with equation  $y = kx$ , where  $k$  is a constant, cuts  $C$  at two distinct points.

(b) Find the range of values for  $k$ .

(6)

$$\begin{aligned} \text{a)} \quad & x^2 + y^2 - 6x + 10y + 9 = 0 \\ \Rightarrow & (x-3)^2 - 9 + (y+5)^2 - 25 + 9 = 0 \\ \Rightarrow & (x-3)^2 + (y+5)^2 = 25 \end{aligned}$$

$$\text{i) centre} = (3, -5)$$

$$\text{ii) radius} = 5$$

b) sub in  $y = kx$

$$\begin{aligned} \Rightarrow & (x-3)^2 + (kx+5)^2 = 25 \\ \Rightarrow & x^2 - 6x + 9 + k^2x^2 + 10kx + 25 = 25 \\ \Rightarrow & x^2(k^2+1) + x(10k-6) + 9 = 0 \end{aligned}$$

$$x = \frac{-(10k-6) \pm \sqrt{(10k-6)^2 - 4 \times (k^2+1) \times 9}}{2(k^2+1)}$$

$$b^2 - 4ac > 0, \quad 100k^2 - 120k + 36 - 36k^2 - 36 > 0$$

$$\Rightarrow 64k^2 - 120k > 0$$

$$k(64k - 120) > 0$$

$$64k > 120$$

$$k > \frac{15}{8}$$

$$\text{So } k > \frac{15}{8} \text{ and } k < 0.$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





15.




Figure 4

Figure 4 shows a sketch of part of the curve  $C$  with equation

$$y = \frac{32}{x^2} + 3x - 8, \quad x > 0$$

The point  $P(4, 6)$  lies on  $C$ .

The line  $l$  is the normal to  $C$  at the point  $P$ .

The region  $R$ , shown shaded in Figure 4, is bounded by the line  $l$ , the curve  $C$ , the line with equation  $x = 2$  and the  $x$ -axis.

Show that the area of  $R$  is 46

(Solutions based entirely on graphical or numerical methods are not acceptable.)

$$y = \frac{32}{x^2} + 3x - 8 \quad (10)$$

$$\frac{dy}{dx} = -2 \frac{32}{x^3} + 3 = -\frac{64}{x^3} + 3$$

gradient of  $C$  at  $P$ : let  $x = 4$ ,  $\frac{dy}{dx} = 2$

Hence gradient of normal ( $l$ ) =  $-\frac{1}{2}$

$$\text{equation of } l: y - 6 = -\frac{1}{2}(x - 4)$$

$$y = -\frac{1}{2}x + 8$$

$$x \text{ intercept of } l: 0 = -\frac{1}{2}x + 8 \quad \text{intercept: } (16, 0).$$

$$x = 16.$$



Question 15 continued

$$\begin{aligned} \text{Area of triangle with points: } (16,0), (4,0) \text{ and } (4,6) \\ = \frac{1}{2} \times 12 \times 6 = 36 \end{aligned}$$

To find area under curve between 2 and 4:

$$\int_2^4 \frac{32}{x^2} + 3x - 8 \, dx$$

$$= \left[ \frac{-1 \cdot 32}{x} + \frac{1}{2} \cdot 3x^2 - 8x \right]_2^4$$

$$= \left( \frac{-32}{4} + \frac{3}{2} \times 4^2 - 32 \right) - \left( \frac{-32}{2} + \frac{3}{2} \times 2^2 - 16 \right)$$

$$= 10$$

$$\text{Total area, } R = 10 + 36 = 46.$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 8 3 4 6 A 0 4 5 4 8